

Physio-biochemical and Enzymatic Responses of Sunflower to Drought Stress

Mozhgan Sarvari¹, Reza Darvishzadeh^{2*}, Roghayeh Najafzadeh² and Hamid Hatami Maleki³

Received: August 14, 2016 Accepted: April 6, 2017

¹Department of Biology, Faculty of Science, Urmia University, Urmia, Iran

²Department of Plant Breeding and Biotechnology, Faculty of Agriculture, Urmia University, Urmia, Iran

³Department of Agronomy and Plant Breeding, Faculty of Agriculture, University of Maragheh, Maragheh, Iran

*Corresponding author; E-mail: r.darvishzadeh@urmia.ac.ir

Abstract

Drought stress is a serious adverse factor that limits sunflower growth and productivity. The stress induces a range of physiological and biochemical responses in sunflower. So, evaluation of defense systems is important for producing resistant cultivars. In this study, physio-biochemical changes and antioxidant enzymes activities of six sunflower lines were evaluated under normal and irrigation at 60 and 40% of the field capacity using a randomized complete block design with three replications. Different characteristics such as relative water content (RWC), chlorophyll a and b contents, carotenoid and proline contents, lipids peroxidation and accumulation of malon dialdehyde (MDA), as well as activity of antioxidative enzymes like guaiacol peroxidase (GPX), ascorbate peroxidase (APX), catalase (CAT) and glutathione reductase (GR) were studied. The results showed significant differences among sunflower lines for physio-biochemical and enzymes activity under drought stress. According to the results, the lines C104 and RHA266 showed the better tolerance to drought stress. The founding of this study can be useful in sunflower breeding programs for producing resistant cultivars to drought stress.

Keywords: Drought tolerance; Enzyme activity; Lipids peroxidation; Physiological characters; Sunflower

Introduction

Water resources are limited and need to be used efficiently in agricultural consumption. Drought stress has been a threat to agricultural activities in many parts of the world (Wang 2004; Vinocur and Altman 2005; Berenguer 2009). The stress affects different morphological, physiological and biochemical characteristics (Reddy *et al.* 2004; Anjum *et al.* 2011; Andrade *et al.* 2013) such as changes in chlorophyll contents (Nayyar and Gupta 2006; Cao *et al.* 2011), inhibition of photochemical activities and decrease the activities of enzymes in the Calvin cycle of photosynthesis (Monakhova and Chernyadev 2002). These changes cause retardation of plant growth and developmental processes. However, the effects of changes are dependent on the time,

stage and severity of water stress (Cao *et al.* 2011). Consequently, the genetic improvement of plant cultivars for water stress tolerance and ability of a crop plant to produce maximum harvest over a wide range of stress and non-stress conditions, has been a principal objective of breeding programs for a long time (Moustafa *et al.* 1996; Chachar *et al.* 2016).

Drought stress induces oxidative stress in plants by generation of reactive oxygen species (ROS) (Farooq *et al.* 2009). The generation of ROS is one of the earliest biochemical responses of cells to the stress. Exceeding of ROS level from removing capacity, leads to oxidative damages including peroxidation of membrane lipids and acclimation of malon dialdehyde (MDA), destruction of photosynthetic pigments

and inactivation of photosynthetic enzymes (Smirnoff 1993). Acclimation of plants to water deficit is the result of different events, which lead to adaptive changes in plant growth and physio-biochemical processes such as changes in plant structure, growth rate, tissue osmotic potential and antioxidant defenses (Duan *et al.* 2007). Plant antioxidant defense systems including enzymatic and non-enzymatic protection have co-evolved with aerobic metabolism to counteract oxidative damage due to ROS (Davar *et al.* 2013). Scavenging of ROS and reducing their damaging effects may correlate with drought tolerance of plants. Efficient scavenging of ROS produced during drought stress requires the action of several enzymatic (APX¹, CAT², SOD³, POX⁴, MDHAR⁵, DHAR⁶, GR⁷) as well as non-enzymatic (ascorbate, phenolic compounds, carotenoids, glutathione, glycine betaine, proline, sugar, polyamines) antioxidants present in tissues (Gill and Tuteja 2010; Karuppanapandian *et al.* 2011).

Sunflower (*Helianthus annuus* L.) is one of the most important oil seed crops (Hussain *et al.* 2015). Although sunflower is moderately tolerant to water stress, its production is greatly affected by drought stress (Pasda and Diepenbrock 1990). Sunflower exhibits a large varietal difference for osmotic adjustment in response to water shortage (Hussain *et al.* 2014). This is proven that sunflower plants exposed to drought is bearing oxidative stress by overproduction of ROS (Rao

2006). Hence, evaluation of enzymatic and non-enzymatic antioxidant defense systems is important in producing resistant sunflower hybrids. Despite several studies concerning drought stress in sunflower (Terbea *et al.* 1995; Baldini *et al.* 1997; Baldini and Vannozzi 1998; Cellier *et al.* 1998; Maury *et al.* 2000; Tahir *et al.* 2002; Poormohammad Kiani *et al.* 2007; Rauf and Sadaqat 2008; Darvishzadeh *et al.* 2010; Safavi *et al.* 2015; Hussein *et al.* 2015), there are fewer reports about simultaneous study of both enzymatic and non-enzymatic antioxidant systems in diverged genotypes of sunflower. The relation between increasing the resistance to drought stress and enhancing the enzymatic antioxidant systems has been studied in some plants species such as rice (Guo *et al.* 2006), sugar beet (Bor *et al.* 2003), wheat (Khanna Chopra and Selote 2007), barley (Acar *et al.* 2001) and sunflower (Ghobadi *et al.* 2013) in abiotic stress conditions. In the present study, some defense systems of sunflower genotypes were studied under drought stress conditions. The information presented here could assist sunflower breeders to choose parents of crosses for breeding of resistance to drought stress.

Materials and Methods

Plant material and experimental methodology

Six oilseed sunflower lines were selected from 125 recombinant inbred lines based on their different responses to drought stress (Poormohammad Kiani *et al.* 2009). Prominent features of the studied lines are summarized in Table 1. The experiment was conducted as the randomized complete block design with three

¹Ascorbate peroxidase (APX)

²Catalase (CAT)

³Superoxide dismutase (SOD)

⁴Peroxidase (POX)

⁵Monodehydroascorbate reductase (MDHAR)

⁶Dehydroascorbate reductase (DHAR)

⁷Glutathione reductase (GR)

replications. The seeds were planted in the plastic pots with a diameter of 12 and height of 14 cm which filled with soil and peat moss (3:1). The plants grew in a greenhouse with 12 h light and maximum and minimum temperature of 28 and 12°C, respectively. Amount of water applied was identical for all treatments from the beginning of planting time until the complete establishment of sunflower plants (eight-leaf (V8) stage). When plants grew into the 8-leaf stage, the control pots were maintained at field capacity and other pots

were irrigated at 60 and 40% of field capacity until the end of growth period. It should be noted that for the drought stress experiment, first the plants were irrigated at 60% of field capacity during 5 days and leaves were sampled on each pot. Then, the pots were irrigated at 40% of field capacity until the end of growth period. Finally, different physio-biochemical parameters and antioxidant enzymes were measured at two levels of drought stress.

Table 1. Characteristics of the studied sunflower lines

| Line | Origin | Type | Characteristics (Poormohammad Kiani <i>et al.</i> , 2007, 2008, 2009) |
|--------|--------|------|---|
| C104 | France | RIL | Good water status and osmotic adjustment as well as biomass and yield under drought stress |
| LR25 | France | RIL | Good water status and osmotic adjustment as well as biomass but it lost grain weight under drought stress |
| LR4 | France | RIL | Average water status and osmotic adjustment as well as biomass and yield under drought stress |
| C100 | France | RIL | Good water status and osmotic adjustment but low in yield under both well-watered and drought stress |
| LR55 | France | RIL | The lowest water status traits and osmotic adjustment as well as biomass and yield under drought stress |
| RHA266 | USA | BL | Low water status traits and osmotic adjustment and average biomass and yield under drought stress |

BL: breeder's line; RIL: recombinant inbred line

Measurement of physio-biochemical characteristics

Relative water content (RWC)

A piece of fresh leaf was removed and fresh weight was measured. The samples were placed in distilled water in a closed container at room temperature for 24 h and then turgid weight was measured. To measure dry weight, the samples were placed in an oven at temperature of 72°C for 24 h and then weighted. Finally, RWC was calculated by the following formula (Levitt 1980):

$$\text{RWC} = (\text{FW} - \text{DW} / \text{TW} - \text{DW}) \times 100$$

In this formula, FW, TW and DW refer to fresh, turgid and dry weight of the leaves, respectively.

Chlorophyll a, b and carotenoid contents

Chlorophyll a, b and carotenoid contents were measured using the method described by Lichtenthaler and Wellburn (1985). An amount of 0.1 g of fresh leaf was homogenized in 5 mL of 100% acetone in a mortar. After homogenizing, the extract was centrifuged for 10 min at 2500 rpm. Then absorbance of the supernatant was recorded at 662, 645 and 470 nm by UV/Vis spectrophotometer (WPA S2100, UK) and the

Chlorophyll a, b and carotenoid contents were calculated using the following formula and expressed in $\mu\text{g/g}$ FW:

$$\text{Chla} = 11.75 A_{662} - 2.350 A_{645}$$

$$\text{Chlb} = 18.61 A_{645} - 3.960 A_{662}$$

$$C_{X+C} = 1000 A_{470} - 2.270 \text{Chla} - 81.4 \text{Chlb} / 227$$

Proline content

Proline content was measured using the method described by Bates *et al.* (1973). An amount of 0.04 g of dried leaf tissue was homogenized in 15 mL of Sulpho salicylic acid (3%) and kept for 72 h in a refrigerator at 4°C to release the proline. After 72 h, the samples were centrifuged at 3000 rpm for 20 min. Then 2 mL of glacial acetic acid and 2 mL of reagent ninhydrin (containing 20 mL phosphoric acid 6 M, 30 mL glacial acetic acid and 1.25 g ninhydrin) were added to 2 mL of the supernatant. The samples were placed in a water bath at 100°C for 1 h. The samples were rapidly cooled using ice and then 4 mL of toluene was added to each sample and stirred. After the formation of two phases, absorption of the supernatant was recorded at 520 nm. To determine the amount of proline, standard curve was prepared using known proline concentrations.

Lipid peroxidation and accumulation of MDA

Lipid peroxidation was measured as an amount of TBARS¹ that determined by TBA reaction described by Heath and Packer (1968). 0.2 g of fresh leaf was homogenized in 5 mL of TCA 1% (w/v) and centrifuged at 8000×g for 10 min. To 1 mL of the supernatant, 4 mL TCA 20% containing TBA 0.5% (w/v) was added. The mixture was

heated at 95°C for 30 min and then quickly cooled on ice. Then the extract was centrifuged at 8000×g for 5 min. The amount of TBARS was determined from the difference between absorbance at 532 and 600 nm using extinction coefficient of 155 $\text{mM}^{-1}\text{cm}^{-1}$ by the following formula:

$$\text{MDA } (\mu\text{mol/gFW}) = (A_{532} - A_{600} / 155) \times 1000$$

Activity of antioxidant enzymes

For extraction of enzymes, 0.5 g of fresh leaf was homogenized in a chilled mortar. Then 3 mL of ice-cold extraction buffer containing Tris-HCl 0.05 M, pH 7.5, MgCl_2 3 mM and EDTA 1 mM was added on powder. The extraction buffer included 2 mM ascorbate that is served for determining APX activity. The homogenate was centrifuged at 15000 g for 15 min at 4°C and the supernatant was served as an enzyme extract. Guaiacol peroxidase (GPX) activity was also determined according to the method of Chang and Kao (1998). The reaction contained 2.5 mL potassium phosphate buffer 50 mM, 1 mL H_2O_2 1% (w/v), 1 mL guaiacol 1% and 0.3 mL extraction buffer. GPX activity was measured at 420 nm. Extinction coefficient of 26.6 ($\text{mM}^{-1}\text{cm}^{-1}$) in a minute was used to calculate its activity. In addition, APX activity was measured using the modified method originally described by Asada (1992). The reaction contained 2.5 mL potassium phosphate buffer (50 mM, pH 7.0) which included ascorbate 0.5 mM and EDTA 0.1 mM. Then, 0.2 mL H_2O_2 1% (w/v) and 0.1 mL extraction buffer were added. After that, APX activity was measured at 240 nm. Extinction coefficient of 2.8 $\text{mM}^{-1}\text{cm}^{-1}$ was used to calculate its activity.

¹Thiobarbituric acid reactive substances (TBARS)

Moreover, CAT activity was determined by measuring H_2O_2 consumption (Maehly and Chance 1959) in a reaction containing 3 mL phosphate buffer (50 mM, pH 7.0), 0.2 mL H_2O_2 1% (w/v) and 50 μL extraction buffer. CAT activity was measured at 240 nm. Extinction coefficient of $0.036 \text{ mM}^{-1} \text{ cm}^{-1}$ was used to calculate its activity. Finally, GR was defined as an amount of the enzyme which decreases A340 (1u per min).

Statistical analysis

The data were analyzed by GLM procedure of SAS 9.2 software. Comparison of mean treatments was made with Tukey test.

Results and Discussion

Relative water content (RWC)

Results pertaining to relative water content revealed significant reduction in RWC of studied lines as compared to the control plants under two drought stresses except for the line LR55 at 60% drought stress condition. Maximum and minimum reduction of RWC was observed in the lines C100

and RHA266, respectively, under two drought stresses (Figure 1a, b). Drought stress resulted in the loss of water in plants as well as reduction of RWC. Therefore, RWC is widely used as one of the most reliable indices for characterization of both sensitivity and tolerance of plants to water stress (Rampino *et al.* 2006). It has been reported that resistant varieties have the highest RWC (Bastide *et al.* 1993; Antolin *et al.* 1993). Rauf and Sadaqat (2008) reported that all the studied sunflower genotypes retained lower RWC under drought regime in comparison to the irrigated condition. Also, Ghobadi *et al.* (2013) observed that RWC decreased in sunflower by drought severity. Significant variability and inhibitory effect has been reported in sunflower for this trait under drought stress condition (Baldini *et al.* 1997). In this study, the more resistant lines had the better osmotic adjustment to prevent their water potential under both drought and normal conditions. The highest RWC was related to the lines LR55 and RHA266 at 60% and the lines C104 and again RHA266 at 40% drought stress.




Figure 1. Relative water content (RWC) of the studied sunflower lines at 60% (a) and 40% (b) drought stress as compared to the normal conditions (100%)

Chlorophyll a, b and carotenoid contents

Chlorophyll a and b contents were increased under both drought stresses as compared to the control except for the line LR55 at 60% drought stress. The highest increase in chlorophyll was observed in the line LR25 at 60% and the line RHA266 at 40% drought stress (Figure 2a, b). Chlorophyll is one of the major chloroplast components for photosynthesis and its content has a positive relationship with photosynthetic rate (Anjum *et al.* 2011). The chlorophyll content is considered as one of the most important indicators of vegetative stage and its degradation is normally considered as a measure of drought susceptibility (Beard 1973; Kim *et al.* 1989; Manivannan *et al.* 2007). In this study, the chlorophyll content decreased with decreasing the irrigation water. The decrease correlates with RWC in the leaves (Munne Bosch and Alegre 2000). Drought stress caused a large decline in the chlorophyll a, chlorophyll b and total chlorophyll content in different sunflower varieties (Manivannan *et al.* 2007). Jabari *et al.* (2009) reported that drought resistant wheat varieties have higher chlorophyll content under drought stress. Ghobadi *et al.* (2013) stated that the chlorophyll a and b content in sunflower decrease with drought severity. Some authors found an opposite trend of

chlorophyll increase by deficit irrigation. Khayatnezhad (2011) and Alaei (2011) showed that drought stress increase leaf chlorophyll content in maize and wheat. In other studies, chlorophyll content was decreased (Jung 2004; Nayyar and Gupta 2006) and increased (Schurr *et al.* 2000; Jiang and Huang 2001; Barraclough and Kate 2001) in plants under drought stress conditions. According to our results, the chlorophyll content increased under drought stress. It seems that increase in chlorophyll content under stress condition, is probably due to the decrease in leaf area which causes accumulation of chlorophyll. The highest chlorophyll content was observed in line LR4 at 60% and line RHA266 at 40% drought stress.

Carotenoid content increased in most of the studied lines as compared to the control under drought stress except for the lines C104, LR25 and LR4 at 60% drought stress. The highest increase in carotenoid content was observed in the line C100 at 60% and the line RHA266 at 40% drought stress (Figure 3a, b). The line C100 had the highest carotenoid content as compared to the others under both drought stress conditions. According to the results of Ghobadi *et al.* (2013), carotenoid content increases in sunflower with drought severity.




Figure 2. Chlorophyll a and b contents of the studied sunflower lines at 60% (a, c) and 40% (b, d) drought stress as compared to the normal conditions (100%)




Figure 3. Carotenoid content of the studied sunflower lines at 60% (a) and 40% (b) drought stress as compared to the normal conditions (100%)

Proline content

There was different response of the genotypes to drought stress in terms of proline content. Based on the Figure 4a, proline level was increased in the lines C104, LR25, LR55 and decreased in the

lines C100, LR4 and RHA266 at 60% drought stress as compared to the control. But at 40% drought stress, the proline content significantly reduced in all of the studied lines, so the maximum and minimum reduction were observed

in the lines C100 and LR25, respectively (Figure 4b). Accumulation of proline occurs in response to the stress (Hare and Cress 1997; Slama *et al.* 2006). Proline acts as a potent scavenger of ROS and prevents the induction of programmed cell death by ROS (Chen and Dickman 2005). Many plants use organic osmolites such as proline for osmotic regulation and to better tolerate the stress. Proline is a reservoir for carbon, nitrogen and also scavenges free radicals. Moreover, proline stabilizes ultra-cellular structures such as membranes and proteins and removes cellular redox potential that is caused by stress (Chen and Muruta 2000; Errabii *et al.* 2006). In many crops such as sorghum, rice, Indian mustard and tomato, differences in proline content has been reported under stress conditions. Ghobadi *et al.* (2013) reported that the proline content in sunflower

increases with drought severity. However, some researchers did not found any increase in proline content under stress condition (Naik and Joshi 1983; Chavan and Karadje 1986). According to our results, the more resistant lines had the highest proline content under both drought and normal conditions. Thus, the highest proline content was seen in lines C104 and LR25 at 60 and 40% drought stress, respectively.

Lipid peroxidation and accumulation of MDA

Regarding data for MDA accumulation that indicates lipid peroxidation, all of the studied lines showed an increase in MDA content as compared to the control under both drought stresses except for the line RHA266 at 60% drought stress. The lowest increase in MDA was observed in the line C100 and LR25 at 60%, and the lines LR25 and LR55 at 40% drought stress




Figure 4. Proline content of the studied sunflower lines at 60% (a) and 40% (b) drought stress as compared to the normal conditions (100%)

(Figure 5a, b). Lipid peroxidation, estimated by the change in MDA content, is used as a reliable marker of oxidative stress (Davar *et al.* 2013). It is reported that the accumulation of MDA under oxidative stress is a by-product of fatty acid

peroxidation, synthesized by cellular membrane lipid peroxidation in different plants species during oxidative stress. Our results are in consistent with the finding of Sairam and Srivastava (2001) that showed the MDA increase

in wheat under drought stress. They have reported that when antioxidant defenses decrease or the formation of free radicals increase, oxidative stress is produced that lead to the increase in the lipid peroxidation of unsaturated fatty acids, lipid membrane damage and subsequent withdrawal of various aldehydes such as MDA. In a study on a salt sensitive variety of wheat, MDA accumulation was greater than the resistant variety (Khanna Chopra and Selote 2007). Our results confirmed that high accumulation of ROS has overcome the antioxidant system of plants and increased membrane lipid peroxidation under stress. Considering these results, the lowest values of MDA belonged to the lines LR4 and RHA266 at 60% drought while the line LR55 possessed lowest values at 40% drought stress.

GPX

As shown in Figure 6a, GPX activity increased significantly in all of the studied lines except for C100 whose activity declined at 60% drought stress. Under 40% drought stress, the GPX activity increased in lines C100, LR55 and RHA266 (Figure 6b). GPX can breaks down indole-3-acetic acid and has important role in lignin biosynthesis and the defense against the live stresses consuming H_2O_2 in cytosol, vacuole, cell wall and the extracellular space (Gill and Tuteja 2010). Increasing of GPX activity has been reported in previous experiments. Smirnov (1993) indicated that increasing of the guaiacol

peroxidase, ascorbate peroxidase and catalase activity is an outcome to the ROS and especially H_2O_2 production under drought stress. Increasing of the GPX activity has also been reported by Zhang *et al.* (1995) in maize under drought stress. Based on the results of Ghobadi *et al.* (2013) POX in sunflower was not affected by drought stress. According to our results, the more resistant and susceptible lines had the highest GPX activity under drought stress. So, the highest GPX activity belonged to lines LR25 and LR55 at 60 and 40% of drought stress, respectively.

APX

APX activity significantly increased in the lines C100 and C104 but, declined in other lines at 60% drought stress (Figure 6c). At the state of 40% drought stress, APX activity significantly increased in the lines LR4 and LR55 (Figure 6d). APX is placed in both the cytosol and chloroplast and can be used to effectively destroy hydrogen peroxide, especially in chloroplasts, where there is no catalase (Grodin and Beck 1979). APX activity was studied under stress conditions and the results showed that in some studies APX activity increased (Sairam *et al.* 2002; Jung 2004), while in some other its activity decreased (Sharma and Dubey 2005). However, in another study no change in APX activity was observed (Bartoli *et al.* 1999). Considering our results, the highest APX activity belonged to the lines C100 and LR4 at 60 and 40% drought stress, respectively.




Figure 5. Malondialdehyde (MDA) content of the studied sunflower lines at 60% (a) and 40% (b) drought stress as compared to the normal conditions (100%)

CAT

CAT activity increased in the lines C100 and RHA266 at 60% drought stress, while it declined significantly in the lines LR4 (Figure 6e). Under 40% drought stress, CAT activity increased significantly in the lines C104, LR55 and RHA266 (Figure 6f). CAT is usually located in peroxisomes and hydrolyzes and detoxifies hydrogen peroxide. CAT activity was studied under stress conditions and the results have shown that in some studies CAT activity increased (Wang *et al.* 2001; Sairam *et al.* 2002), while in some other its activity decreased (Jiang and Huang 2001; Sharma and Dubey 2005). However, in another study no change in APX activity was observed (Bartoli *et al.* 1999). Ghobadi *et al.* (2013) reported that the CAT activity increases in sunflower with drought severity. Also, Jung (2004) showed that the CAT and POX activity in the mature leaves of tall grass increased as compared to the control under water stress conditions. But any increase was not observed in the young leaves. In another study, it was observed that the activity of these two enzymes

increase during the early stages of water stress. It seems that water stress prevents the detrimental effects of ROS on cell membrane via increasing the CAT activity as much as possible (Jiang and Huang 2001). Considering our results, CAT activity increased in the line RHA266 under both water stress conditions compared to the control. The highest CAT activity observed in the line C100 at 60% and in the lines C104 and RHA266 at 40% drought stress.

GR

GR activity increased in the lines C100 and RHA266 at 60% drought stress and the increase was higher in RHA266 (Figure 6g). Under 40% drought stress, level of the GR activity increased in the lines C100, C104, LR4 and RHA266 while it decreased in the lines LR25 and LR55 (Figure 6h). GR is the last enzyme in glutathione-ascorbate pathway which consumes NADPH as an electron donor to reduce glutathione (Noctor and Foyer 1998). Increasing of GR activity in cells is a way to increase the glutathione level. This increases the cell tolerance toward ROS




Figure 6. Antioxidant enzymes' activity of the studied sunflower lines at 60% (a, c, e, g) and 40% (b, d, f, h) drought stress as compared to the normal conditions (100%)

(Jiang and Huang 2001). Regarding the increase in cellular GR activity under drought stress and its role in glutathione reduction, GR is probably one of the most important enzymes in plant that the increase in its activity can enhance plant tolerance against oxidative stress (Khanna Chopra and Selote 2007). There are several conflicting reports about the activity of the GR under drought stress. According to the reports, both increasing (Sairam *et al.* 2002; Jung 2004) and decreasing (Jiang and Huang 2001) of the GR activity have been reported under drought stress. Considering our results, the highest GR activity belonged to the lines C100 and RHA266 at 60% and the line C104 at 40% drought stress.

Conclusion

The results showed significant differences for physio-biochemical and enzymatic responses of

sunflower lines to the drought stress. Resistant and susceptible lines showed different responses to the stress. Some lines which were recognized as a drought sensitive genotype based on non-enzymatic variables were drought resistant genotype based on enzymatic characters. However, the antioxidant enzymes' activity was much more pronounced in the resistant lines of sunflower compared to the susceptible lines. It seems that each of enzymatic and non-enzymatic variables could clarify some aspects of drought resistance and must be used simultaneously for the screening of sunflower genotypes against drought stress. The lines C104 and RHA266 had good water status and osmotic adjustment and showed better tolerance to the drought stress. The finding of this study can be useful in sunflower breeding programs for producing resistant cultivars to drought stress.

References

- Acar O, Turkan I and Zdemir FO, 2001. Superoxide dismutase and peroxidase activities in drought sensitive and resistant barley (*Hordeum vulgare* L.) varieties. *Acta Physiologiae Plantarum* 3: 351-356.
- Alaei Y, 2011. The effect of amino acids on leaf chlorophyll content in bread wheat genotypes under drought stress conditions. *Middle-East Journal of Scientific Research* 10: 99-101.
- Andrade A, Vigliocco A, Alemano S, Llanes A and Abdala G, 2013. Comparative morpho-biochemical responses of sunflower lines sensitive and tolerant to water stress. *American Journal of Plant Sciences* 4: 156-167.
- Anjum SH, Xie XY, Wang LC, Saleem MF, Man C and Lei W, 2011. Morphological, physiological and biochemical responses of plants to drought stress. *African Journal of Agricultural Research* 6: 2026-2032.
- Antolin MC and Sanchez-Diaz M, 1993. Effect of temporary drought on photosynthesis of alfalfa plants. *Journal of Experimental Botany* 44: 1341-1349.
- Asada RD, 1992. Ascorbate peroxidase—a hydrogen peroxide scavenging enzyme in plants. *Plant Physiology* 85: 235-241.
- Baldini M and Vannozzi GP, 1998. Agronomic and physiological assessment of genotypic variation for drought tolerance in sunflower genotypes obtained from a cross between *H. annuus* and *H. argophyllus*. *Agricultural Medicine* 128: 232-240.
- Baldini M, Vannozzi GP, Gomez Sanchez D and Turi M, 1997. Results of a physiological selection approach to improve drought resistance in sunflower. *Genetika-a-Slechtenti* 33: 181-195.
- Barracough PB and Kyte J, 2001. Effect of water stress on chlorophyll meter reading in wheat. *Plant Nutrition* 92: 722-23.

- Bartoli CG, Simontacchi M, Tambussi E, Beltrano J, Montaldi E and Puntarulo S, 1999. Drought and watering-dependent oxidative stress. Effect on antioxidant content in *Triticum aestivum* L. leaves. *Journal of Experimental Botany* 332: 375-383.
- Bastide B, Deborah S, Janet H and Lrwin T, 1933. Effects of severe water stress on aspects of crassulacean acid metabolism in *Xerosicyos dangui*. *Plant Physiology* 103: 1089-1096.
- Bates LS, Walderon RP and Teare ID, 1973. Rapid determination of free proline for water stress studies. *Plant and Soil* 39: 205-208.
- Beard JB, 1973. *Turf Grass: Science and Culture*. Prentice-Hall, Englewood Cliffs, NJ, USA.
- Bor M, Ozdemir F and Turkan I, 2003. The effect of salt stress on lipid peroxidation and antioxidants in leaves of sugar beet *Beta vulgaris* L. and wild beet *Beta maritima* L. *Plant Science* 164: 77-84.
- Cao HX, Sun CX, Shao HB and Lei XT, 2011. Effects of low temperature and drought on the physiological and growth changes in oil palm seedlings. *African Journal of Biotechnology* 10: 2630-2637.
- Cellier F, Conejero G, Breitler JC and Casse F, 1998. Molecular and physiological responses to water deficit in drought-tolerant and drought-sensitive lines of sunflower: accumulation of dehydrin transcripts correlates with tolerance. *Plant Physiology* 116: 319-328.
- Chachar MH, Chachar NA, Chachar Q, Mujtaba SM, Chachar S and Chachar Z, 2016. Physiological characterization of six wheat genotypes for drought tolerance. *International Journal of Research-Granthaalayah* 4: 184-196.
- Chang CJ and Kao CH, 1998. H₂O₂ metabolism during senescence of rice leaves: change in enzyme activities in light and darkness. *Plant Growth Regulation* 25: 11-15.
- Chavan PD and Karadge BA, 1986. Growth, mineral nutrition, organic constituents and rate of photosynthesis in *Sesbania grandiflora* L. grown under saline conditions. *Plant and Soil* 93: 395-404.
- Chen C and Dickman MB, 2005. Proline suppresses apoptosis in the fungal pathogen *Colletotrichum trifolii*. *Proceedings of the National Academy of Sciences* 102: 3459-3464.
- Chen THH and Murata N, 2000. Enhancement of tolerance of abiotic stress by metabolic engineering of betaines and other compatible solutes. *Current Opinion in Plant Biology* 5: 250-257.
- Darvishzadeh R, Pirzad A, Hatami Maleki H, Poormohammad Kiani S and Sarrafi A, 2010. Evaluation of the reaction of sunflower inbred lines and their F1 hybrids to drought conditions using various stress tolerance indices. *Spanish Journal of Agricultural Research* 8: 1037-1046.
- Davar R, Darvishzadeh R and Majd A, 2013. Changes in antioxidant systems in sunflower partial resistant and susceptible lines as affected by *Sclerotinia sclerotiorum* (Lib.) de Bary. *Biologia* 68: 821-829.
- Duan B, Yang Y, Lu Y, Korpelainen H, Berninger F and Li C, 2007. Interactions between drought stress, ABA and genotypes in *Picea asperata*. *Journal of Experimental Botany* 58: 3025-3036.
- Errabii T, Gandonou CB, Essalmani H, Abrini J, Idaomar M and Skalisenhaji N, 2006. Growth, proline and ion accumulation in sugarcane callus cultures under drought induced osmotic stress and its subsequent relief. *African Journal of Biotechnology* 5: 1488-1493.
- Farooq M, Wahid A, Kobayashi N, Fujita D and Basra SMA, 2009. Plant drought stress: effects, mechanisms and management. *Agronomy for Sustainable Development* 29: 185-212.
- Ghobadi M, Shayesteh Taherabadi SH, Ghobadi ME, Mohammadi GR and Honarman SJ, 2013. Antioxidant capacity, photosynthetic characteristics and water relations of sunflower (*Helianthus annuus* L.) cultivars in response to drought stress. *Industrial Crops and Products* 50: 29-38.
- Gill SS and Tuteja N, 2010. Reactive oxygen species and antioxidant machinery in abiotic stress tolerance in crop plants. *Plant Physiology and Biochemistry* 48: 909-930.
- Groden D and Beck E, 1979. H₂O₂ destruction by ascorbate-dependent system from chloroplast. *Biochimica et Biophysica Acta* 546: 426-435.
- Guo Z, Ou W, Lu S and Zhong Q, 2006. Differential responses of antioxidative system to chilling and drought in four rice cultivars differing in sensitivity. *Plant Physiology and Biochemistry* 44: 828-836.
- Hare PD and Cress WA, 1997. Metabolic implications of stress induced proline accumulation in plants. *Plant Growth Regulation* 21: 79-102.
- Heath RL and Packer L, 1968. Photoperoxidation in isolated chloroplasts. *Archives of Biochemistry and Biophysics* 125: 850-857.
- Hussain S, Saleem MF, Iqbal J, Ibrahim M, Ahmad M, Nadeem SM, Ali A and Atta S, 2015. Abscisic acid mediated biochemical changes in sunflower (*Helianthus annuus* L.) grown under drought and well-watered field conditions. *The Journal of Animal and Plant Sciences* 25: 406-416.

- Hussain S, Saleem MF, Iqbal J, Ibrahim M, Atta S, Ahmed T and Rehmani MIA, 2014. Exogenous application of abscisic acid may improve the growth and yield of sunflower hybrids under drought. *Pakistan Journal of Agricultural Sciences* 51: 49-58.
- Jabari F, Ahmadi A, Poustini K, Alizadeh H, Sharifzadeh F and Ranjbar M, 2009. Evaluation of relationship between relative water content and gas exchanges parameters with drought resistance in 7 wheat cultivars. *Iranian Journal of Field Crop Science* 40: 198-207 (In Persian with the English abstract).
- Jiang Y and Huang N, 2001. Drought and heat stress injury to two cool-season turf grass in relation to antioxidant metabolism and lipid peroxidation. *Crop Science* 41: 436-442.
- Jung S, 2004. Variation in antioxidant metabolism of young and mature leaves of *Arabidopsis thaliana* subjected to drought plant. *Plant Sciences* 166: 459-466.
- Karuppanapandian T, Moon JH, Kim C, Manoharan, K and Kim W, 2011. Reactive oxygen species in plants: their generation, signal transduction and scavenging mechanisms. *Australian Journal of Crop Science* 5: 709-725.
- Khanna Chopra R and Selote DS, 2007. Acclimation to drought stress generates oxidative stress tolerance in drought resistant than susceptible wheat cultivar under field conditions. *Environmental and Experimental Botany* 60: 276-283.
- Khayatnezhad M, Gholamin RS, Jamaatie Somarin H and Zabihie Mahmoodabad R, 2011. The leaf chlorophyll content and stress resistance relationship considering in corn cultivars (*Zea mays* L.). *Advances in Environmental Biology* 5: 118-122.
- Kim KS, Beard JB and Sifers SI, 1989. Drought resistance comparisons among major warm-season turf grasses. *USGA Green Section Record* 26(5): 12-15.
- Levitt J, 1980. *Response of Plants to Environmental Stresses: Water, Radiation, Salt and Other Stresses*. Academic Press, New York.
- Lichtenthaler HK and Wellburn AR, 1985. Determination of total carotenoids and chlorophylls a and b of leaf in different solvents. *Biochemical Society Transactions* 11: 591-592.
- Lopez-Berenguer C, Martinez Ballesta MC, Moreno DA, Carvajal M and Garcia Viguera C, 2009. Growing hardier crops for better health: salinity tolerance and the nutritional value of broccoli. *Journal of Agricultural and Food Chemistry* 57: 572-578.
- Maehly AC and Chance B, 1959. The assay of catalase and peroxidase. In: Click D (Eds). *Method of Biochemical Analysis*. Pp. 357-425. Interscience Publishers, New York.
- Manivannan P, Jaleel CA, Sankar B, Kishorekumar A, Somasundaram R, Alagu Lakshmanan GM and Panneerselvam R, 2007. Growth, biochemical modifications and proline metabolism in *Helianthus annuus* L. as induced by drought stress. *Colloids and Surfaces B: Biointerfaces* 59: 141-149.
- Maury P, Berger M, Mojayad F and Planchon C, 2000. Leaf water characteristics and drought acclimation in sunflower genotypes. *Plant and Soil* 223: 153-160.
- Monakhova OF and Chernyadev II, 2002. Protective role of kartolin-4 in wheat plants exposed to soil drought. *Applied and Environmental Microbiology* 38: 373-380.
- Moustafa MA, Boersma L and Kronstad WE, 1996. Response of four spring wheat cultivars to drought stress. *Crop Science* 36: 982-986.
- Munne Bosch S and Alegre L, 2000. Changes in carotenoids, tocopherols and diterpenes during drought and recovery, and the biological significance of chlorophyll loss in *Rosmarinus officinalis* plants. *Planta* 210: 925-931.
- Naik CR and Joshi CL, 1983. Ineffectual role of proline metabolism in salt stressed sugarcane leaves. *Proceedings of the Indian Academy of Science* 92: 265-269.
- Nayyar H and Gupta D, 2006. Differential sensitivity of C₃ and C₄ plant to water deficit stress: association with oxidative stress and antioxidants. *Environmental and Experimental Botany* 58: 106-113.
- Noctor G and Foyer C, 1998. Ascorbate and glutathione: keeping active oxygen under control. *Plant Physiology* 49: 249-279.
- Pasda G and Diepenbrock W, 1990. The physiological yield analysis of sunflower. Part II. Climate factors. *Wissenschaft Technologie* 93: 155-168.
- Poormohammad Kiani S, Grieu P, Maury P, Hewezi T, Gentzbittel L and Sarrafi A, 2007. Genetic variability for physiological traits under drought conditions and differential expression of water stress-associated genes in sunflower (*Helianthus annuus* L.). *Theoretical and Applied Genetics* 114: 193-207.

- Poormohammad Kiani S, Maury P, Sarrafi A and Grieu P, 2008. QTL analysis of chlorophyll fluorescence parameters in sunflower (*Helianthus annuus* L.) under well-watered and water-stressed conditions. *Plant Science* 175: 565-573.
- Poormohammad Kiani S, Nouri L, Maury P, Darvishzadeh R, Grieu P and Sarrafi A, 2009. Genetic variation and identification of molecular markers associated with osmotic adjustment-related traits in gamma irradiation-induced mutants of sunflower (*Helianthus annuus* L.). *Journal of Genetics and Breeding* 62: 67-74.
- Rampino P, Pataleo S, Gerardi C, Mita G and Perrotta C, 2006. Drought stress response in wheat: physiological and molecular analysis of resistant and sensitive genotypes. *Plant Cell and Environment* 29: 2143-2152.
- Rao KVM, 2006. Introduction. In: Rao KVM, Raghavendra AS and Reddy KJ (Eds). *Physiology and Molecular Biology of Stress Tolerance in Plants*. Pp.1-14. Springer, The Netherlands.
- Rauf S and Sadaqat HA, 2008. Identification of physiological traits and genotypes combined to high achene yield in sunflower (*Helianthus annuus* L.) under contrasting water regimes. *Australian Journal of Crop Science* 1: 23-30.
- Reddy AR, Chaitanya KV and Vivekanandan M, 2004. Drought induced responses of photosynthesis and antioxidant metabolism in higher plants. *Journal of Plant Physiology* 161: 1189-1202.
- Safavi SM, Safavi AS and Safavi SA, 2015. Evaluation of drought tolerance in sunflower (*Helianthus annuus* L.) inbred lines and synthetic varieties under non Stress and drought stress conditions. *Biological Forum* 17: 1849-1854.
- Sairam R, Rao K and Srivastava C, 2002. Differential response of wheat genotypes to long term salinity stress in relation to oxidative stress, antioxidant activity and osmolyte concentration. *Plant Science* 163: 1037-1046.
- Sairam RK and Srivastava C, 2001. Water stress tolerance of wheat (*Triticum aestivum* L): variations in hydrogen peroxide accumulation and antioxidant activity in tolerant and susceptible genotypes. *Journal of Agronomy and Crop Science* 186: 63-70.
- Schurr U, Heckenberger U, Herdel K, Walter A and Feil R, 2000. Leaf development in *Ricinus communis* during drought stress: dynamics of growth processes of cellular structure and of sink-source transition. *Journal of Experimental Botany* 51: 1515-1529.
- Sharma P and Dubaey RS, 2005. Drought induces oxidative stress and enhances the activities of antioxidant enzymes in growing rice seedling. *Plant Growth Regulation* 46: 209-221.
- Slama I, Messedi D, Ghnaya T, Savoure A and Abdely C, 2006. Effects of water deficit on growth and proline metabolism in *Sesuvium portulacastrum*. *Environmental and Experimental Botany* 56: 231-238.
- Smirnoff N, 1993. The role of active oxygen in the response of plant to water deficit and desiccation. *New Phytologist* 125: 27-58.
- Tahir MHN, Muhammad I and Hussain MK, 2002. Evaluation of sunflower (*Helianthus annuus* L.) inbred lines for drought tolerance. *International Journal of Agriculture and Biology* 4: 398-400.
- Terbea M, Vranceanu AV, Petcu E, Craiciu DS and Micut G, 1995. Physiological response of sunflower plants to drought. *Romanian Agricultural Research* 3: 61-67.
- Vinocur B and Altman A, 2005. Recent advances in engineering plant tolerance to abiotic stress: achievements and limitations. *Current Opinion in Biotechnology* 16: 123-132.
- Wang JR, Li SX and Li KI, 2001. Effect of water limited deficit stress during different growth stages on leaf enzymes of winter wheat. *Acta Botanica Boreali-Occidentalia Sinica* 21: 47-52.
- Wang L, Liu D, Ahmed T, Chung FL, Conaway C and Chiao JW, 2004. Targeting cell cycle machinery as a molecular mechanism of sulforaphane in prostate cancer prevention. *International Journal of Oncology* 24: 187-192.
- Zhang J, Cui SLIJ and Kirkham MB, 1995. Protoplasmic factors, antioxidant responses, and chilling resistance in maize. *Plant Physiology and Biochemistry* 33: 567-575.