

Methyl Jasmonate-Induced Changes in Non- and Antioxidant-Enzymatic Defense in Peppermint (*Mentha piperita*)

Soheila Afkar¹, Ghasem Karimzadeh^{1*}, Mokhtar Jalali Javaran¹, Mozafar Sharifi² and Mehrdad Behmanesh³

Received: January 26, 2013 Accepted: May 11, 2013

¹Department of Plant Breeding and Biotechnology, Faculty of Agriculture, Tarbiat Modares University, Tehran, Iran

²Plant Biology Department, Faculty of Biological Sciences, Tarbiat Modares University, Tehran, Iran

³Genetics Department, Faculty of Biological Sciences, Tarbiat Modares University, Tehran, Iran

*Corresponding author Email: karimzadeh_g@modares.ac.ir

Abstract

Peppermint (*Mentha piperita* L.), a herbaceous and perennial species which is produced mainly for the medicine and food. The peppermint plants were initiated from 10 cm-long rhizome cuttings followed by transferring into pots. The 48 h-treated plants with methyl jasmonate (MJ) concentrations (0, 0.1, 0.5 mM) were assessed for their total soluble proteins, malondialdehyde (MDA), chlorophylls a, b and total, anthocyanin, total carbohydrates, carotenoid, activity of antioxidant guaiacol peroxidase (POD) and superoxide dismutase (SOD) enzymes. The data were analyzed using completely randomized design (CRD) with three replications. Mean comparisons were carried out, using Duncan's multiple range test. MJ treatment caused significant changes in soluble proteins, chlorophylls (a, b and total), MDA, carbohydrates and antioxidant enzymes (SOD and POD) but had no effect on anthocyanin and carotenoid. These results indicate that MJ can effectively improve the defense system and antioxidant capacity of peppermint.

Keywords: Anthocyanin; Antioxidant enzymes; Chlorophyll; Malondialdehyde; Methyl jasmonate; Peppermint

Abbreviations CAR–Carotenoid; CAT–Catalase; Chl–Chlorophyll; CHO–Carbohydrate; H₂O₂–Hydrogen peroxide; JA–Jasmonic acid; JAs–Jasmonates; MDA–Malondialdehyde; MJ–Methyl jasmonate; POD–Guaiacol peroxidase; ROS–Reactive oxygen species; SOD–Superoxide dismutase

Introduction

Peppermint (*Mentha piperita* L.) belongs to mint (Lamiaceae) family and is considered as a medicinal and aromatic plant species. Peppermint essential oil includes menthol, menthone, methylacetat, menthofuran and pulegone (Mahmoud and Croteau 2003; Tabatabaie and Nazar 2007). Its cultivation has economic importance, due to its ability to produce and store essential oil, whose main constituent is menthol, used in oral hygiene products, pharmaceuticals, cosmetics and foods. Menthol also has high antifungal and antibacterial potentials, thus becoming one of the most demanded substances

by the scents and essences industry (Scavroni 2005). Because of this and other reasons, peppermint essential oil ranks high in terms of total sales volume (Orozco-Cárdenas *et al.* 2001). This herb synthesizes and concentrates oils in its leaves in highly specialized epidermal secretory structures known as glandular trichomes (McCaskill *et al.* 1992). Jasmonates (JAs) including jasmonic acid (JA) and MJ are a family of cyclopentanone compounds synthesized from linolenic acid via the octadecanoic pathway. They inhibit plant growth generally but also promote diverse processes as a class of plant growth regulator consisting of fruit ripening, senescence,

tuber formation, tendril coiling, pollen formation and defense-related responses against mechanical and insect wounding and pathogen infection (Ueda and Kato 1980; Creelman and Mullet 1997). The jasmonates applied exogenously to plants exert various effects either inhibiting or promoting the morphological and physiological changes. It has been shown that MJ causes the generation of H_2O_2 (Orozco-Cárdenas and Ryan 1999; Orozco-Cárdenas *et al.* 2001; Hung and Kao 2004) and lipid peroxidation expressed as MDA production in plant cells (Hung and Kao 1998, 2004). Thus, MJ leads to oxidative stress in plant cells. Plants have an internal protective enzyme catalyzed clean up system to scavenge reactive oxygen species (ROS), thus ensuring normal cellular function. Superoxide dismutase (SOD) constitutes the first line of defense via detoxification of super oxide radicals (Sairam and Saxena 2000), thereby maintaining membranes of plant tissue. SOD detoxifies superoxide anion free radicals by forming H_2O_2 ; It can be further eliminated by concerted action of catalase (CAT) and POD. In addition, MJ helps in maintaining the pools of antioxidant enzymes and alleviating the oxidative stress (Li *et al.* 1998; Jung 2004). Both SOD and POD are important enzymes associated with anti-oxidative stress in plants. ROS scavenging group depends on the detoxification mechanism provided by an integrated system of non-enzymatic reduced molecules and enzymatic antioxidants (Jaleel *et al.* 2006). Exogenously applied JA and MJ lead to decreased expression of photosynthesis-related genes encoding for example the small subunit of ribulose-1,5-bisphosphate carboxylase/oxygenase (Rubisco),

reduced translation and increased degradation of Rubisco and rapid loss of chlorophyll (Chl) in barley leaves (Weidhase *et al.* 1987; Parthier 1990). The MJ does not only regulate a variety of plant-developmental responses, but is also induced by pathogen attack or wounding, which often leads to the generation of ROS, including H_2O_2 , superoxide anions ($O_2^{\cdot-}$) and hydroxyl free radicals (OH) (Faurie *et al.* 2009; Parra-Lobato *et al.* 2009). ROS have the potential to interact with many cellular components, triggering stresses in plant cell culture, leading to membrane damage and, as a result, there is an immediate cellular response to trigger plant defense signals. Plants possess antioxidant defense systems, consisting of enzymatic and non-enzymatic components, which normally maintain ROS balance within the cell. Plants contain substantial amounts of carotenoids that serve as non-enzymatic scavengers of ROS (Young and Britton 1990). Anti-oxidative enzymes include SOD, which catalyzes the disproportion of superoxide radicals to hydrogen peroxide and POD, which removes H_2O_2 (Kumari *et al.* 2006). The POD is associated with biochemical and physiological processes such as growth, cell formation, fruit development, ethylene biosynthesis, as well as the response to various stresses (Matamoros *et al.* 2003).

The purpose of this study was to determine the changes in the physiological characteristics and in the activities of antioxidant enzymes capacity in *Mentha piperita* treated with different concentrations of MJ. It was hypothesized that MJ could improve non- and antioxidant enzymatic-defense in peppermint.

Materials and Methods

This experiment was carried out in the greenhouse of Tarbiat Modares University, Tehran, Iran. The peppermint plants were supplied from Iranian Institute of Medicinal Plants, Karaj, Iran. They initiated from 10 cm-long rhizome cuttings followed by transferring into pots. The 48 h-treated plants with MJ on three different concentrations (0, 0.1, 0.5 mM) were assessed for their total soluble proteins, chlorophylls (a, b, and total), MDA, total carbohydrates, carotenoid, anthocyanin and antioxidant enzymes (SOD and POD).

Determination of protein, lipid peroxides, carbohydrates and chlorophyll content in leaf extract

Soluble protein extraction was carried out according to Ausubel *et al.* (1995) and determined with Folin-Ciocalteu reagent according to Lowry *et al.* (1951) and Bradford (1976). The level of lipid peroxidation was measured in terms of MDA content, a product of lipid peroxidation, following the method of De Vos (1991). MDA is a major cytotoxic product of lipid peroxidation and acts as an indicator of free radical production. Total carbohydrates were estimated spectrophotometrically according to the method of Dubois *et al.* (1956). Chlorophyll was extracted in 80% (v/v) acetone from the leaf samples according to the method of Arnon (1949).

Measurement of carotenoid, anthocyanin content, POD and SOD activity

Carotenoid and anthocyanin were estimated spectrophotometrically according to the methods

of Helrich (1990) and Krizek *et al.* (1993), respectively. POD activity was determined as an increase in optical density due to the formation of guaiacol dehydrogenation product according to Kar and Mishra (1976). SOD activity was assayed by using the photochemical NBT following the method of Giannopolitis and Ries (1977).

Statistical analysis

The data were analyzed using completely randomized design with three replications by Minitab 16. Means and standard errors (SE) were used to compare MJ treatments, using Duncan's multiple range test. Moreover, correlation coefficients were calculated among all physiological characteristics.

Results

The result of ANOVA showed that MJ had significant effect on most measured physiological characteristics in the leaves of peppermint (Table 1). Antioxidant reactions in MJ-treated *Mentha piperita* caused a significant decrement in photosynthetic activities and pigment levels. The contents of Chls a, b and total decreased significantly in 0.1 mM MJ-treated leaves (Figures 1A, B, C, respectively) compared with the control, but no significant changes in those characters were detectable in 0.5 mM MJ-treated leaves. The ratio of Chls a + b/CAR also decreased remarkably in the MJ treated leaves (Figure 1D), indicating that the changes of total Chls a + b takes place faster than that of total carotenoids (CARs). In the present study, the MDA concentration increased significantly when plants were subjected to 0.1 mM MJ treatment

compared with the control. CAR content was not changed by inducing MJ treatment, while protein content decreased in 0.5 mM MJ-treated leaves (Figure 1F). Total carbohydrates (CHO; glucose) content in 0.5 mM MJ-induced leaves was detected to be lower than in those of controls (Figure 1G). Our data showed that treatment of *Mentha piperita* plants with 0.5 mM MJ leads to a

significant increase in POD activity. Total POD and SOD activities showed prominently about 2.36- and 1.81-fold increase, respectively at 48 h of MJ induction (Figures 1H, I). Antioxidant enzymes exhibited the highest activities at 0.5 mM MJ exposure compared to those in the control plants.

Table 1. Effect of MJ on physiological characters in *Mentha piperita*

SOV	df	MS										
		Protein	Antocyanin	Chl a	Chl b	Total Chl	CAR	MDA	POD	SOD	Total CHO	Chl a +b/CAR
MJ	2	4.53 ^{***}	0.00008 ^{ns}	2.728 ^{**}	3.564 ^{**}	2.275 [*]	0.0169 ^{ns}	0.156 ^{**}	2.995 ^{**}	2.995 ^{**}	102.94 ^{**}	2.728 ^{**}
Error	6	0.18	0.00004	0.258	0.182	0.409	0.0053	0.013	0.169	0.169	9.43	0.258

^{ns}, ^{*}, ^{**} and ^{***} Non significant and significant at 5%, 1% and 0.1% probability levels, respectively

Correlation coefficients among physiological characters (Table 2) showed positive and highly significant relationship between POD and SOD ($r = 0.985^{***}$) and negative and highly significant correlations between soluble protein and either SOD ($r = -0.911^{**}$) or POD ($r = -0.918^{***}$).

Similarly, total CHO negatively and significantly correlated with either Chl a ($r = -0.781^*$), Chl b ($r = -0.745^*$) or total Chl ($r = -0.704^*$). There were positive and significant correlations between total CHO and either protein ($r = 0.727^*$) or MDA ($r = 0.847^{**}$).

Table 2. Correlation coefficients among physiological characters in *Mentha piperita*

	Protein	Antocyanin	Chl a	Chl b	Total Chl	CAR	MDA	POD
Antocyanin	-0.189 ^{ns}							
Chl a	-0.511 ^{ns}	-0.336 ^{ns}						
Chl b	-0.398 ^{ns}	-0.398 ^{ns}	0.964 ^{***}					
Total Chl	-0.391 ^{ns}	-0.413 ^{ns}	0.979 ^{***}	0.985 ^{***}				
CAR	-0.792 [*]	0.124 ^{ns}	0.521 ^{ns}	0.456 ^{ns}	0.461 ^{ns}			
MDA	0.713 [*]	0.150 ^{ns}	-0.786	-0.777 [*]	-0.778 [*]	-0.666 [*]		
POD	-0.918 ^{***}	0.280 ^{ns}	0.445 ^{ns}	0.318 ^{ns}	0.295 ^{ns}	0.583 ^{ns}	0.576 ^{ns}	
SOD	-0.911 ^{**}	0.256 ^{ns}	0.461 ^{ns}	0.353 ^{ns}	0.336 ^{ns}	0.585 ^{ns}	-0.664 ^{ns}	
Total CHO	0.727 [*]	0.174 ^{ns}	-0.781 [*]	-0.745 [*]	-0.704 [*]	-0.638 ^{ns}	0.847 ^{**}	
Chl a+b/CAR	0.074 ^{ns}	-0.494 ^{ns}	0.867 ^{**}	0.904 ^{**}	0.918 ^{***}	0.075 ^{ns}	-0.568 ^{ns}	-0.497 ^{ns}

^{ns}, ^{*}, ^{**} and ^{***} Non significant and significant at 5%, 1% and 0.1% probability levels, respectively

Discussion

Hormones inclusive jasmonates may mediate the response of plants to environmental stresses and may interact with other cellular metabolites and environmental factors in the regulation of stress responses (Parthier 1990). In previous reports, it was found that MJ stimulates the production of H_2O_2 (Orozco-Cárdenas and Ryan 1999, Orozco-Cárdenas *et al.* 2001; Hung and Kao 2004), leading to oxidative stress in plant cells. Reduced oxygen species such as hydrogen peroxide (H_2O_2) and superoxide radicals (O^{2-}) are formed due to oxidative stress and they can produce free radicals, inducing lipid peroxidation and protein denaturation. The MDA is an oxidized product of membrane lipids and its level can show the extent of oxidative stress. When plants were treated with 0.1 mM MJ, the MDA concentration significantly increased compared to the control. Smaller amount of MDA by 0.5 mM MJ application in our study (Figure 1E) indicated that it had better efficiency to endure the damage of cellular membranes than 0.1 mM MJ concentration. MDA, produced by lipid peroxidation of cell membrane, is often used as an indicator of salt and oxidative damages (Mandhania 2006).

Decrease in MDA level by 0.5 mM MJ application may be the result of increased activities of antioxidant enzymes that can help to clean up ROS and alter ratio of membranes fatty acids as a major source of ROS production (Wang 1999). However, lipid peroxidation operated under exogenous application of MJ in peanut (Kumari *et al.* 2006). Previous studies showed soluble protein content as a main index of physiological condition of plants. We can express

the disturbance in protein metabolism as a reason for decreasing the total soluble protein amount of the treated plants (Figure 1F). Many results suggested a connection between photosynthesis and jasmonates in plants. The contents of Chls a, b and total decreased significantly in 0.1 mM MJ-treated leaves (Figures 1A, B, C, respectively) compared with the control. It was reported that Chl a is more intensely degraded than Chl b (Wolf 1956). Exogenously applied MJ reduced translation and increased degradation of Rubisco and resulted in rapid loss of Chl in barley leaves (Weidhase *et al.* 1987; Parthier 1990). Results of the present study indicated that significant increase in total CHO is somehow associated with reduction in Chl. Sugar, mainly glucose, accumulation in the cell is responsible for the regulation of photosynthetic process (Moore *et al.* 1999). The highly significant correlations among CHO, protein and MDA (Table 2) suggested that physiological traits have a close relationship with each other. The important components of thylakoid membranes are CARs which can effectively suppress the excited Chl a, Chl b and total Chl (Knox and Dodge 1985).

Further increment in antioxidant enzyme activities is caused by exogenous application of MJ (Anjum *et al.* 2011)., The modification of antioxidant enzymes (SOD, POD) can play important protective roles in avoiding the deleterious effects triggered by elevated levels of ROS observed at initial moments of MJ exposure. To minimize the damaging results of ROS, plants use a lot of evolved non- and enzymatic-antioxidant systems. Plants contain substantial amounts of CAR that serve as non-enzymatic




Figure 1. Effects of methyl jasmonate (MJ) on A) Chl a (mg g⁻¹ F.W.), B) Chl b (mg g⁻¹ F.W.), C) Total Chl (mg g⁻¹ F.W.), D) Chl a+b/CAR, E) MDA (mM cm⁻¹), F) protein (mg g⁻¹ F.W.), G) total CHO (mg g⁻¹FW), H) POD (Δ A₄₇₀ mg⁻¹ protein) and I) SOD (mg mg⁻¹ protein) in *Mentha piperita*.

M0 = Control

M1 = 0.1 mM

M2 = 0.5

scavengers of ROS (Young and Britton 1990). The metabolism of ROS is dependent on several functionally interrelated antioxidant enzymes such as SOD and POD. Enzymatic antioxidant systems provide protection against the toxic effects of ROS (Scandalios 1993). PODs are involved in a large number of biochemical and

physiological processes (Yip 1964) and may change quantitatively and qualitatively during growth and development (Shannon 1969). The SOD is believed to play a crucial role in antioxidant defense because it catalyzes the dismutation of O²⁻ into H₂O₂, whereas CAT and POD destroy H₂O₂ (Scandalios 1993). A positive

and high correlation between POD and SOD (Table 2) suggests that an increase of SOD activity was accompanied by an increase of POD activity as a result of high demand of quenching H₂O₂. Such findings in *Mentha piperita* are consensus with *Triticum aestivum* results reported by Ghobadi *et al.* (2011). In *Arabidopsis*, when JA was used at a concentration of 100 µM, no significant alteration in the enzymes activities was detected until the third day of induction, but the activity was reduced remarkably after 6 d of treatment (Berger 2002). In barley, MJ mediated the stimulation of antioxidant enzymes including SOD, CAT and POD (Popova *et al.* 2003). JA ability to cause chlorosis led to the suggestion that this compound plays a role in plant senescence (Ueda *et al.* 1981), however, it was reputed by the fact that high JA levels were found in the zones of cell division, young leaves, and reproductive structures (Creelman and Mullet 1997). Application of MJ caused a senescence-like symptom as indicated by a great decline in photosynthesis and Chls and a strong increase in anthocyanins and antioxidant enzyme activities in *Arabidopsis thaliana* (Jung 2004). The most obvious character of leaf senescence is yellowing. Chl loss has been the principal criterion of senescence in the most reports. The protein degradation during leaf senescence has been

realized in the earliest studies. In the present study, the senescence of *Mentha piperita* leaves was followed by measuring the decrease of Chl and protein contents. It is clear that MJ significantly promotes the senescence of peppermint leaves. These results are in agreement with those in the previous reports (Chao and Kao 1992; Tsai *et al.* 1996; Chen and Kao 1998). Several reports showed that the soluble sugar content often goes up, not down, in senescing leaves (Shiroya *et al.* 1961; Trippi 1965; Egli *et al.* 1980; Lazan *et al.* 1983; Crafts-Brandner *et al.* 1984). It has even been proposed that elevated sugar content actually causes senescence (Lazan *et al.* 1983). Our results showed that glucose (soluble sugar) content decreased in 0.5 mM MJ-treated peppermint leaves (Figure 1G), refuting the suggestion that sugar accumulation may cause leaf senescence. Moreover, the effect of exogenous MJ treatment on antioxidant enzymes (POD, SOD) and non-enzymatic defenses was evaluated, verifying that MJ can increase the activity of antioxidant enzymes in *Mentha piperita*.

Acknowledgments

Authors would like to thank Tarbiat Modares University for financially supporting this research work.

References

- Anjum SA, Wang L, Farooq M, Khan I and Xue L, 2011. Methyl jasmonate-induced alteration in lipid peroxidation, antioxidative defence system and yield in soybean under drought. *Journal of Agronomy and Crop Science* 197(4): 296-301.
- Arnon DI, 1949. Copper enzymes in isolated chloroplasts, polyphenoloxidase in *Beta vulgaris*. *Plant Physiology* 24: 1-15.
- Ausubel FM, Brent R and Kingston RE (Eds.), 1995. *Current Protocols in Molecular Biology*. John Wiley and Sons, New York, USA.

- Berger S, 2002. Jasmonate-related mutants of *Arabidopsis* as tools for studying stress signaling. *Planta* 214: 497-504.
- Bradford M, 1976. A rapid and sensitive method for quantitation of microgram quantities of protein utilizing the principle of "protein- dye binding". *Annual Review of Biochemistry* 72: 248-254.
- Chao CM and Kao CH, 1992. Methyl jasmonate, calcium and leaf senescence in rice. *Plant Physiology* 99: 1693-1694.
- Chen SJ and Kao CH, 1998. Methyl jasmonate, ammonium and leaf senescence in rice. *Journal of Plant Physiology* 152: 353-357.
- Crafts-Brandner SJ, Below FE, Wittenbach VA, Harper, JR and Hageman RH, 1984. Differential senescence of maize hybrids following ear removal. II. Selected leaf. *Plant Physiology* 74: 368-373.
- Creelman RA and Mullet JE, 1997. Biosynthesis and action of jasmonates in plants. *Annual Review of Plant Physiology* 48: 355-381.
- De Vos C, Schat HM, De Waal MA, Vooijs R and Ernst W, 1991. Increased to copper-induced damage of the root plasma membrane in copper tolerant *Silene cucubalus*. *Plant Physiology* 82: 523-528.
- Dubois M, Gilles KA, Hamilton JK, Rebers PA and Smith F, 1956. Colorimetric method for determination of sugars and related substances. *Analytical Chemistry* 28: 350-356.
- Egli DB, Laggett JE and Cheniae A, 1980. Carbohydrate levels in soybean leaves during reproductive growth. *Crop Science* 20: 468-473.
- Faurie B, Cluzet S and Merillon JM, 2009. Implication of signaling pathways involving calcium, phosphorylation and active oxygen species in methyl jasmonate-induced defense responses in grapevine cell cultures. *Journal of Plant Physiology* 166: 1863-1877.
- Ghobadi M, Khosravi S, Kahrizi D and Shirvani F, 2011. Study of water relations, chlorophyll and their correlation with grain yield in wheat (*Triticum aestivum* L.) genotypes. *World Academy Science, Engineering and Technology* 78: 582-585.
- Giannopolitis CN and Ries SK, 1977. Superoxide dismutases: II. Purification and quantitative relationship with water-soluble protein in seedlings. *Plant Physiology* 59: 315- 318.
- Helrich K (Ed), 1990. Official methods of analysis, 15th edn. Association of Official Analytical Chemistry. Washington, DC, pp. 17-33.
- Hung KT and Kao CH, 1998. Involvement of lipid peroxidation in methyl jasmonate-promoted senescence in detached rice leaves. *Plant Growth Regulation* 24: 17-21.
- Hung KT and Kao CH, 2004. Nitric oxide acts as an antioxidant and delays methyl jasmonate-induced senescence of rice leaves. *Journal of Plant Physiology* 161: 43-52.
- Jaleel CA, Gopi R, Manivannan P, Kishorekumar A, Sankar B and Panneerselvam R, 2006. Paclobutrazol influences on vegetative growth and floral characteristics of *Catharanthus roseus* (L.) G. Don. *Indian Journal of Applied and Pure Biology* 21: 369-372.
- Jung S, 2004. Effect of chlorophyll reduction in *Arabidopsis thaliana* by methyl jasmonate or norflurazon on antioxidant systems. *Journal of Plant Physiology and Biochemistry* 42: 231-255.
- Kar M and Mishra D, 1976. Catalase, peroxidase and polyphenol-oxidase activities during rice leaf senescence. *Plant Physiology* 57: 315-319.
- Knox JP and Dodge AD, 1985. Singlet oxygen and plants. *Phytochemistry* 24: 889-896.
- Krizek DT, Kramer GF, Upadhyaya A and Mirecki RM, 1993. UV-B response of cucumber seedling grown under metal halide and high pressure sodium/deluxe lamps. *Plant Physiology* 88: 350-358.
- Kumari G, Reddy A, Naik S, Kumar S, Prasanthi J, Sriranganayakulu G, Reddy P and Sudhakar C, 2006. Jasmonic acid induced changes in protein pattern, antioxidative. *Plant Biology* 50 (2): 219-226.
- Lazan HB, Barlow EW and Bardy CJ, 1983. The significance of vascular connection in regulating senescence of the detached flag leaf of wheat. *Journal of Experimental Botany* 34: 726-736.
- Li L, Staden JV and Jager AK, 1998. Effect of plant growth regulators on the antioxidant system in seedlings of two maize cultivars subjected to water stress. *Journal of Plant Growth Regulation* 25: 81-87.
- Lowry OH, Rosebrough NT, Farr AL and Randall RJ, 1951. Protein measurement with the folin phenol reagent. *Journal of Biological Chemistry* 193: 265-275.
- Mahmoud SS and Croteau RB, 2003. Menthofuran regulates essential oil biosynthesis in peppermint by controlling a downstream monoterpene reductase. *Proceedings of the National Academy of Sciences* 100 (24): 14481-14486.
- Mandhanian S, Madan S and Sawhney V, 2006. Antioxidant defense mechanism under salt stress in wheat seedlings. *Plant Biology* 50: 227-231.
- Matamoros MA, Dalton DA, Ramos J, Clemente MR, Rubio MC and Becana M, 2003. Biochemistry and molecular biology of antioxidants in the rhizobialegume symbiosis. *Plant Physiology* 133: 499-509.
- McCaskill D, Gersheuzon, J and Croteau R, 1992. Morphology and monoterpene biosynthetic capabilities of secretory cell clusters isolated from glandular trichomes of peppermint (*Mentha piperita* L.). *Planta* 187: 445-454.
- Moore DB, Cheng SH, Sims D and Seemann JR, 1999. The biochemical and molecular basis for photosynthetic acclimation to elevated atmospheric CO₂. *Plant Cell and Environment* 22: 567-582.

- Orozco-Cárdenas ML, Narva'ez-Vázquez J and Ryan CA, 2001. Hydrogen peroxide acts as a second messenger for the induction of defense genes in tomato plants in response to wounding, systemin and methyl jasmonate. *Plant Cell* 13: 179-191.
- Orozco-Cárdenas ML and Ryan CA, 1999. Hydrogen peroxide is generated systemically in plant leaves by wounding and systemin via the octadecanoid pathway. *Proceedings of the National Academy of Sciences* 96: 6553-6557.
- Parra-Lobato MC, Fernandez-Garcia N, Olmos E, Alvarez-Tinaut MC and Gomez-Jimenez MC, 2009. Methyl jasmonate-induced antioxidant defence in root apoplast from sunflower seedlings. *Environmental and Experimental Botany* 66: 9-17.
- Parthier B, 1990. Jasmonates: hormonal regulators or stress factors in leaf senescence. *Journal of Plant Growth Regulation* 9 (1): 57-63.
- Popova L, Ananieva E, Hristova V, Christov K, Georgieva K, Alexieva V and Stoinova ZH, 2003. Salicylic acid and methyl jasmonate-induced protection on photosynthesis to paraquat oxidative stress. *Bulgarian Journal of Plant Physiology* 29 (3-4): 133-152.
- Sairam RK and Saxena DC, 2000. Oxidative stress and antioxidants in wheat genotypes: possible mechanism of water stress tolerance. *Journal of Agronomy and Crop Science* 184 (1): 55-61.
- Scandalios JG, 1993. Oxygen stress and superoxide dismutases. *Plant Physiology* 101: 7-12.
- Scavroni J, Boaro CSF, Ortiz M, Marques M and Leonardo CF, 2005. Yield and composition of the essential oil of *Mentha piperita* L. (Lamiaceae) grown with biosolid. *Brazilian Journal of Plant Physiology* 17 (4): 345-352.
- Shannon LM, 1969. Plant isoenzymes. *Annual Review of Biochemistry* 38: 189-210.
- Shiroya M, Lister GR, Nelson VD and Krotkov G, 1961. Translocation of ^{14}C in tobacco at different stages of development following assimilation of $^{14}\text{CO}_2$ by a single leaf. *Canadian Journal of Botany* 39: 855-864.
- Tabatabaie J and Nazar J, 2007. Influence of nutrient concentrations and NaCl salinity on the growth, photosynthesis and essential oil content of peppermint and *Lemon verbena*. *Turkish Journal of Agriculture and Forestry* 31: 245-253.
- Trippi VS, 1965. Studies on ontogeny and senility in plants. XI. Leaf shape and longevity in relation to photoperiodism in *Gaillardia pulchella*. *Phyton* 22: 113-117.
- Tsai FY, Hung KT and Kao CH, 1996. An increase in ethylene sensitivity is associated with jasmonate-promoted senescence of detached rice leaves. *Journal of Plant Growth Regulation* 15: 197-200.
- Ueda J and Kato J, 1980. Identification of a senescence-promoting substance from wormwood (*Artemisia absinthum* L.). *Plant Physiology* 66: 246-249.
- Ueda J, Kato J, Yamane H and Takahashi N, 1981. Inhibitory effect of methyl jasmonate and its related compounds on kinetin-induced retardation of oat leaf senescence. *Plant Physiology* 52: 305-309.
- Wang SY, 1999. Methyl jasmonate reduces water stress in strawberry. *Journal of Plant Growth Regulation* 18: 127-134.
- Weidhase RAE, Kramell HM, Lehmann J, Liebisch HW, Lerbs W and Parthier B, 1987. Methyl jasmonate-induced changes in the polypeptide pattern of senescing barley leaf segments. *Plant Science* 51: 177-186.
- Wolf FT, 1956. Changes in chlorophylls a and b in autumn leaves. *American Journal of Botany* 43: 714-718.
- Yip CC, 1964. The hydroxylation of proline by horseradish peroxidase. *Biochemica et Biophysica Acta* 92: 395-396.
- Young A and Britton G, 1990. Carotenoids and stress, In: Alscher RG and Cummings JR (Eds.). *Stress Responses in Plants: Adaptation and Acclimation Mechanisms*, pp. 87-112. Wiley-Liss Inc., New York, USA.