

Supplementary Table 2. Shoot regeneration from internodal stems and pulvini of petiole explants cultured on the shoot induction medium with BAP and IBA combinations.

PGRs (mg/L)		Internodal stem		Pulvinus of the petiole	
BAP	IBA	Transverse TCL	Longitudinal TCL	Transverse TCL	Longitudinal TCL
0	0	0.0±0.0k	0.0±0.0k	0.0±0.0k	0.0±0.0k
0.25	0	0.90±0.003j	1.1±0.003j	0.9±0.003j	1.20±0.003j
0.5	0	4.25±0.012h	4.73±0.013h	4.55±0.13h	5.30±0.015h
0.75	0	7.9±0.022e	7.88±0.022g	7.8±0.022f	8.1±0.023f
1.0	0	2.75±0.00i	3.15±0.009i	3.6±0.01i	3.90±0.011i
0	0.25	0.0±0.0k	0.0±0.0k	0.0±0.0k	0.0±0.0k
0.25	0.25	0.0±0.0k	0.0±0.0k	0.0±0.0k	0.0±0.0k
0.5	0.25	10.03±0.03c	10.35±0.03d	10.75±0.031c	10.70±0.031e
0.75	0.25	10.95±0.032b	10.97±0.03c	11.70±0.034b	11.40±0.032d
1.0	0.25	6.25±0.02f	5.93±0.017g	6.0±0.02g	6.20±0.018g
0	0.5	0.0±0.0k	0.0±0.0k	0.0±0.0k	0.0±0.0k
0.25	0.5	0.0±0.0k	0.0±0.0k	0.0±0.0k	0.0±0.0k
0.5	0.5	0.0±0.0k	0.0±0.0k	0.0±0.0k	0.0±0.0k
0.75	0.5	21.52±0.065a	14.77±0.07a	26.3±0.064a	24.90±0.071a
1.0	0.5	9.25±0.02d	13.25±0.038b	9.8±0.03d	13.40±0.038b
0	0.75	0.0±0.0k	0.0±0.0k	0.0±0.0k	0.0±0.0k
0.25	0.75	0.0±0.0k	0.0±0.0k	0.0±0.0k	0.0±0.0k
0.5	0.75	0.0±0.0k	0.0±0.0k	0.0±0.0k	0.0±0.0k
0.75	0.75	0.0±0.0k	0.0±0.0k	0.0±0.0k	0.0±0.0k
1.0	0.75	4.35±0.012g	9.65±0.027e	9.3±0.026e	12.60±0.036c

The results of four completely randomized designs are expressed as means ± SE; Means with different letters in each column are significantly different at $p \leq 0.05$ according to Tukey's test; PGRs: Plant growth regulators; TCL: Thin cell layer; Measurements were taken after eight weeks of culture.