

**Supplementary Table 4.** Shoot regeneration from internodal stems and pulvini of petiole explants cultured on the shoot induction medium with BAP and NAA combinations.

PGRs (mg/L)		Internodal stem		Pulvinus of the petiole	
		BAP	NAA	Transverse TCL	Longitudinal TCL
0	0	0.0±0.0k		0.0±0.0k	0.0±0.0k
0.25	0	0.90±0.003j		0.90±0.003j	0.90±0.003j
0.5	0	4.25±0.012b		4.25±0.012b	4.25±0.012b
0.75	0	<b>7.9±0.022a</b>		<b>7.9±0.022a</b>	<b>7.9±0.022a</b>
1.0	0	2.75±0.00c		2.75±0.00c	2.75±0.00c
0	0.25	0.0±0.0k		0.0±0.0k	0.0±0.0k
0.25	0.25	0.0±0.0k		0.0±0.0k	0.0±0.0k
0.5	0.25	1.8±0.005g		1.8±0.005g	1.8±0.005g
0.75	0.25	2.18±0.006f		2.18±0.006f	2.18±0.006f
1.0	0.25	1.53±0.004h		1.53±0.004h	1.53±0.004h
0	0.5	0.0±0.0k		0.0±0.0k	0.0±0.0k
0.25	0.5	0.0±0.0k		0.0±0.0k	0.0±0.0k
0.5	0.5	0.0±0.0k		0.0±0.0k	0.0±0.0k
<b>0.75</b>	<b>0.5</b>	2.4±0.007d		2.4±0.007d	2.4±0.007d
1.0	0.5	2.28±0.006e		2.28±0.006e	2.28±0.006e
0	0.75	0.0±0.0k		0.0±0.0k	0.0±0.0k
0.25	0.75	0.0±0.0k		0.0±0.0k	0.0±0.0k
0.5	0.75	0.0±0.0k		0.0±0.0k	0.0±0.0k
0.75	0.75	0.0±0.0k		0.0±0.0k	0.0±0.0k
1.0	0.75	1.13±0.003i		1.13±0.003i	1.13±0.003i

PGRs: Plant growth regulators; TCL: Thin cell layer; The results of four completely randomized designs are expressed as means ± SE; Means with different letters in each column are significantly different at  $p \leq 0.05$  according to Tukey's test; Measurements were taken after eight weeks of culture.